

## PHYTOCHEMICAL ANALYSIS OF PLANT SPECIES OF GENUS ZANTHOXYLUM

N.W. AYANGLA<sup>1</sup>, NEETU SINGH<sup>2</sup> & AJAY KUMAR<sup>3</sup>

<sup>1</sup>Department of Biotechnology, Mewar University, Gangrar, Chittorgarh, Rajasthan, India

<sup>2,3</sup>Department of Biotechnology, Mewar Institute of Management, Ghaziabad, Uttar Pradesh, India

### ABSTRACT

The present article illustrates phytochemical investigations carried out on three plant species i.e. *Zanthoxylum armatum* (D.C), *Zanthoxylum oxyphyllum* (Edgew) and *Zanthoxylum rhetsa* (Roxb.) (D.C). Percent yield of the crude ethanolic extracts from *Zanthoxylum armatum* (Leaves), *Zanthoxylum oxyphyllum* (Leaves), *Zanthoxylum oxyphyllum* (Seeds), *Zanthoxylum rhetsa* (Leaves), *Zanthoxylum rhetsa* (Seeds) were 7.78%, 9.85%, 14.1%, 9.23%, and 14.21% respectively. The qualitative analysis showed the abundance of Glycosides, Coumarins, Flavonoids, Phenols, and Tannins in crude ethanolic extracts of ZAL, ZOL, ZOS, ZRL, and ZRS. Thus, the preliminary phytochemical screening revealed the presence of medicinally important constituents in the plants studied. Phenolic content was found maximum in leaves of *Zanthoxylum oxyphyllum* followed by leaves of *Zanthoxylum armatum*. Seeds of *Zanthoxylum rhetsa* showed maximum concentration of Flavonoid followed by leaves of *Zanthoxylum armatum*. The present qualitative and quantitative analysis explores the wider possibility of efficient and productive extraction of different secondary metabolites from crude ethanolic extracts of ZAL, ZOL, ZOS, ZRL, and ZRS.

**KEYWORDS:** Phytochemical Screening, *Zanthoxylum armatum* (D.C), *Zanthoxylum oxyphyllum* (Edgew) and *Zanthoxylum rhetsa* (Roxb.) (D.C)

**Received:** Nov 29, 2015; **Accepted:** Dec 15, 2015; **Published:** Dec 16, 2015; **Paper Id.:** IJMPSFEB20161

### INTRODUCTION

The potential of plants to serve human beings in a range of aspects has been well documented since antiquity (Albuquerque *et al.*, 2011; Qureshi *et al.*, 2014; Sangeetha and Baskar, 2015). Plant products have been part of Phytomedicines since time immemorial (Criagg and David, 2001). A/c to Wadood *et al.*, (2013), presence of phytochemicals is considered as active medicinal chemical constituents. Medicinal plants play a vital role in preventing various diseases (Wadood *et al.*, 2013) and their medicinal value lies in some bioactive constituents (like flavonoids and phenols) that produce a definite physiological action on the human body (Mir *et al.*, 2013). Phenols have antibacterial, antifungal, nematicidal, insecticidal, antioxidant, antimutagenic, and other activities (Oksana *et al.*, 2012).

A/c to Brijwal *et al.*, (2013), India has richest plant based medicinal traditional system because of its rich biodiversity. Exploration of medicinal plants in the North east India is a matter of attraction to the scientific fertility, traders as well as pharmaceuticals (Shankar and Rawat, 2013). Chakraborty *et al.*, (2012) emphasized on the need of detailed and systematic review on ethanobotanical studies in the North East India. Many species of *Zanthoxylum* are used in traditional medicine especially in America, Africa, and Asia (Prieto *et al.*, 2010). Species of *Zanthoxylum* Genus possess high medicinal, economical as well as ecological importance (Brijwal *et al.*, 2013)

coupled with significant insecticidal and antifungal activities and it is a promising source of various secondary metabolites (Patino *et al.* 2012). Therefore, efforts were made to characterize the different types of metabolites presented in selected species of *Zanthoxylum* Genus followed by their quantitative estimation based on preliminary phytochemical screening.

## RESEARCH METHODOLOGY

Leaves of *Zanthoxylum armatum* (ZAL), *Zanthoxylum oxyphyllum* (ZOL), *Zanthoxylum rhetsa* (ZRL) and Seeds of *Zanthoxylum oxyphyllum* (ZOS), *Zanthoxylum rhetsa* (ZRS) were collected from Chungliyimsen village, District-Mokokchung, Nagaland (North-East India). Plant materials were cleansed with distilled water and allowed to dry for 15 days in dark. Thereafter, it was coarsely crushed using homogenizer and grinded mechanically of mesh size 1 mm. The powdered plant material was extracted successively with 70% Ethanol. After 48 hours, extracts were filtered by using muslin (cheese cloth) followed by Whatman filter paper No.1 and filtrates were evaporated to dryness and weighed. The crude extracts were stored in air tight glass containers at 4°C till further analysis. All extracts were subsequently stored at -20 °C in deep freezer until bioassays were conducted. Solvents used in extraction and Reagents for phytochemical analysis was of pure analytical grade.

Phytochemical investigations of crude alcoholic extracts of ZAL, ZOL, ZRL, ZOS, and ZRS include: Percent of yield of Extract (Patil and Gaikwad, 2010); Preliminary Phytochemical screening (Harbone, 1998; Fransworth, 1996; Rangari, 2002; Khandelwal, 2005; Sofowara, 1993; Trease and Evans, 1989; Harborne, (1973); Total Phenolic Content (Singleton *et al.*, 1999) and Total Flavonoid Content (Nguyen and Eun, 2011).

## RESULTS AND DISCUSSIONS

### Percentage Yield of Extract

Percent yield of the crude ethanolic extracts from *Zanthoxylum armatum* (Leaves), *Zanthoxylum oxyphyllum* (Leaves), *Zanthoxylum oxyphyllum* (Seeds), *Zanthoxylum rhetsa* (Leaves), *Zanthoxylum rhetsa* (Seeds) were 7.78%, 9.85%, 14.1%, 9.23%, and 14.21% respectively. Highest yield was observed in seeds of *Zanthoxylum rhetsa* followed by seeds of *Zanthoxylum oxyphyllum*. Similar results were also reported in the studies conducted on different species of *Zanthoxylum* using different plant parts. 1.45% of extract yield was observed by Islam *et al.*, (2014) in 95% crude ethanolic extract of *Zanthoxylum budrunga* seeds. Reddy and Jose (2011) reported 1.83% of extract yield from aqueous extract of *Zanthoxylum rhetsa* seeds, while Buragohain *et al.*, (2011) reported 1.22 % of extract yield in methanolic extract of *Zanthoxylum oxyphyllum* tender shoots. Present findings were well correlated with the findings of Verma and Khosa (2012).


### Preliminary Phytochemical Analysis

Phytochemical screening is a preliminary step in the characterization of plant products. Preliminary screening for presence or absence of a particular class of compound is qualitative in nature and form a robust base for the quantitative estimation of bioactive components. In the present study, phytochemical components in crude ethanolic extracts of ZAL, ZOL, ZOS, ZRL, and ZRS were investigated through a series of chemical tests. Phytochemicals were screened for Alkaloids, Carbohydrates, Glycosides, Coumarins, Proteins & Amino acids, Phytosterols, Phenolic compounds, Tannins, Flavonoids, Anthocyanins, Lignins, Terpenoids, Phlobatannins and Quinones.

The preliminary phytochemical screening had showed the abundance of Glycosides, Coumarins, Flavonoids, Phenols, and Tannins in crude ethanolic extracts of ZAL, ZOL, ZOS, ZRL, and ZRS. However, Phytosterols, Anthocyanins and Phlobatannins were not detected in all the tested samples. It was observed that the ZOS and ZRS showed abundance of Flavonoids and Coumarins followed by Phenols and Tannins. ZAL, ZOL and ZRL showed significant presence of Tannins followed by Glycosides. Crude ethanolic extract of leaves of *Zanthoxylum armatum* showed abundance of Flavonoids, Glycosides, Tannins and traces of Phenols, Carbohydrates, Coumarins and Lignins. Ethanolic extract of ZOL was screened positive for Phenols, Tannins, Coumarins and Glycosides. Considerable presence of Phenols, Flavonoids and Coumarins followed by Alkaloids, Carbohydrates, and Tannins with traces of Glycosides and Quinones were screened in the crude ethanolic extract of ZOS. Maximum number of secondary metabolites was screened in crude ethanolic extract of ZRS with abundance of Flavonoids, Glycosides, Coumarins, Proteins & amino acids, Tannins followed by Phenols, Alkaloids, Terpenoids, with traces of Carbohydrates and Quinones. Crude ethanolic extract of ZRL was found positive in preliminary screening for Glycosides, Phenols, Flavonoids, Coumarins and Tannins. The present qualitative analysis explores the wider possibilities for efficient and productive extraction of different secondary metabolites from crude ethanolic extracts of ZAL, ZOL, ZOS, ZRL, and ZRS.

### Estimation of Secondary Metabolites

Plant phenolics are the major group of chemical species that act as primary antioxidants (Hatano *et al.*, 1989) and have high redox potentials which allow them to act as oxygen quenchers (Kahkonen *et al.*, 1999) and are responsible for multiple biological effects (Pise *et al.*, 2010).



**Figure 1: Comparative Analysis of Total Phenol and Flavonoid Content in Selected Plant Parts of *Zanthoxylum* Species Using Ethanol as a Solvent**

- *Zanthoxylum armatum* (Leaves); 2) *Zanthoxylum oxyphyllum* (Leaves); 3) *Zanthoxylum oxyphyllum* (Seeds); 4) *Zanthoxylum rhetsa* (Leaves); 5) *Zanthoxylum rhetsa* (Seeds).

Figure 1 shows variation in Total Phenol and Flavonoid content. Phenolic content was found maximum in ethanolic extract of *Zanthoxylum oxyphyllum* leaves followed by *Zanthoxylum armatum* leaves. However, minimum concentration of Phenol was observed in ethanolic extract of *Zanthoxylum rhetsa* leaves. Ethanolic extract of *Zanthoxylum rhetsa* seeds showed maximum concentration of Flavonoids followed by the leaves of *Zanthoxylum armatum*. While analysing the trend in Phenol and Flavonoid content in the leaves and seeds of *Zanthoxylum oxyphyllum*, it was observed

that the ethanolic extract of their leaves were showed higher concentration in comparison to seeds. However, in case of *Zanthoxylum rhetsa*, the trend was opposite. In this case, it was observed that the content of Phenol and Flavonoid was higher in ethanolic extract of seeds instead of leaves. Ethanolic extract of *Zanthoxylum oxyphyllum* leaves analysed with higher Phenolic and Flavonoid content in comparison to their seeds. Previous studies also reported significant yield in Total Phenol and Flavonoid contents from crude extracts of *Zanthoxylum* species. Prabhash *et al.*, (2014) found 62.3 µg/ml of Total Phenol from methanolic extracts of *Zanthoxylum rhetsa* Leaves. A/c to Mahadkar *et al.*, (2013), *Zanthoxylum rhetsa* fruits with methanolic extracts yields  $0.061 \pm 0.29$  g/100g FW of Phenols. *Zanthoxylum rhetsa* bark with ethyl acetate and butanol fractions yields  $20.47 \pm 0.09$  and  $14.14 \pm 0.185$  mg GAE/g DW of Phenols, respectively (Santhanam *et al.*, 2013). Vidyamadhavi *et al.*, (2014) found the Flavonoid yield of 258 µg, 533 µg and 349 µg equivalent of tannic acid (Vidyamadhavi *et al.*, 2014) in ethyl acetate, methanol and aqueous extracts of aerial parts of *Zanthoxylum rhetsa*. *Zanthoxylum armatum* leaves extract yields 28.46 mg/L gallic acid equivalent of Phenols (Mahadkar *et al.*, 2013). Statistical analysis was also employed in order to estimate the correlation between Phenol and Flavonoid content among the selected plant parts (Leaves or Seeds) of *Zanthoxylum armatum*, *Zanthoxylum oxyphyllum*, and *Zanthoxylum rhetsa* using ethanol as a solvent. The strongest correlation was observed in total phenolic and total flavonoid content of *Zanthoxylum oxyphyllum* and *Zanthoxylum rhetsa* leaves and seeds.

## CONCLUSIONS

Traditional medicine is pervasively used by nearly 80% of the population in African countries (WHO, 2000). In China, 40% of health care depends on traditional medicine (WHO, 1999). In India there are about 45,000 plant species, out of them several thousands have been claimed to possess medicinal properties (Grover *et al.*, 2002). A/c to the MOEF (GOI) report, the herbal industry in India uses about 8000 medicinal plants (Gundimeda *et al.*, 2006; Kumar and Janagam, 2011). Another literature estimates that in India there are 880 medicinal plants species involved in all India trade out of which about 48 species are exported (EXIM; Gundimeda *et al.*, 2006; Kumar and Janagam, 2011). The international market turnover for herbal products is of \$6.2 billion; however India's share in the global medicinal plants related export trade is just 0.5 percent (WHO, 2002; Kumar and Janagam, 2011). In order to materialize the biodiversity rich potential, India must focus and develop the scientific cultivation, post harvest technology, processing, manufacturing, patenting, marketing, etc., for medicinal plants, as suggested by Singh, (2006).

A/c to Wadood *et al.*, (2013), the phytochemical analysis of the medicinally important plants has commercial interest in both research institutions and pharmaceutical organizations (Wadood *et al.*, 2013). The present phytochemical study investigates qualitative screening and quantitative analysis of *Zanthoxylum armatum* (D.C), *Zanthoxylum oxyphyllum* (Edgew) and *Zanthoxylum rhetsa* (Roxb.) (D.C) plants. Preliminary Phytochemical screening showed the presence of Alkaloids, Carbohydrates, Glycosides, Coumarins, Proteins & Amino acids, Phenolic compounds, Tannins, Flavonoids, Lignins, Terpenoids and Quinones in crude ethanolic extracts of ZAL, ZOL, ZOS, ZRL and ZRS. The results of phytochemical analysis revealed the presence of medicinally important constituents in the plant species studied. The present study strongly recommended the further work to isolate, purify, characterize, and standardize the bioactive constituents responsible for the medicinal properties of plants belongs to Genus *Zanthoxylum*.

## ACKNOWLEDGEMENTS

We acknowledge all the ones who dedicated themselves in this study by any means. Errors, if any, are purely unintentional. We attempted to trace and acknowledge the copyright holders of all materials reproduced in this paper and to copyright holders if permission and acknowledgements to publish in this form have not been given.

## REFERENCES

1. Albuquerque et al., (2011). Allelopathy, an alternative tool to improve cropping systems-A Review. *Agron. Sustain. Dev.*, 31: 379-395.
2. Brijwal, L., Pandey, A. and Tamta, S. (2013). An overview on phytomedicinal approaches of *Zanthoxylum armatum* DC: An important magical medicinal plant. *Journal of Medicinal Plants Research*, 7(8): 366-370.
3. Buragohain, J., Konwar, B.K. and Bordoloi, M.J. (2011). Isolation of an antimicrobial compounds from the tender shoots of *Zanthoxylum*. *Der Pharmacia Sinicia*, 2(6): 149.
4. Chakraborty, R., Deb, B., Devannac, N. and Sena, S. (2012). North-East India an Ethnic Storehouse of Unexplored Medicinal Plants. *J. Nat. Prod. Plant Resour.*, 2(1): 143-152.
5. Cragg G.M. and David, J.N. (2001). Natural product drug discovery in the next millennium. *Journal of Pharmaceutical Biology*, 39: 8-17.
6. EXIM. (2003). Export potential of Indian medicinal plants and products, Export-Import Bank of India, Occasional Paper no 98. Mumbai: Quest Publications.
7. Farnsworth, N.R. (1996). Biological and phytochemical screening of plants. *J. pharm.Sci*, 55: 225-276.
8. Grover, J.K., Yadav, S. and Vats V. (2002). Medicinal plants of India with anti-diabetic potential. *J Ethnopharmacol.*, 81(1):81-100.
9. Gundimeda, H., Sanyal, S., Sinha, R. and Sukhdev, P. (2006). The value of biodiversity in India's forests. Monograph-4: Green Accounting for Indian States and Union Territories Project. Published by TERI Press, New Delhi for Green Indian States Trust, India, pp:51.
10. Harborne, J.B. (1973). *Phytochemicals Methods*. Chapman and Hall Ltd., London, pp: 49-188.
11. Harborne, J.B. (1998). *Phytochemical methods*. 3rd Edition London: Chapman and Hall.
12. Hatano, T., Edmatsu, R., Mori, A., Fujita, Y. and Yasuhara, E. (1989). Effects of tannins and related polyphenols on superoxide anion radical and on 1, 1-diphenyl-2-picrylhydrazyl radical. *Chem. Pharm Bull.*, 37: 2016-2023.
13. Islam et al., (2014). Antinociceptive and Antioxidant Activity of *Zanthoxylum budrunga* Wall (Rutaceae) Seeds. *The Scientific World Journal*, Vol. 2014, Article ID 869537, pp:7. <http://dx.doi.org/10.1155/2014/869537>.
14. Khandelwal, K.R. (2005). *Practical Pharmacognosy, Techniques and Experiments*. Nirali Prakashan (Pune), 13<sup>th</sup> Edn.
15. Kumar, M.R. and Janagam, D. (2011). Export and import pattern of medicinal plants in India. *Indian Journal of Science and Technology*, 4(3): 245-248.
16. Mahadkar, S., Jadhav, V. and Deshmukh, S. (2013). Antioxidant activity of some promising wild edible fruits. *Der Chemica Sinica*, 4(3): 165-169.
17. Mir, M.A., Sawhney, S.S. and Jassal, M.M.S. (2013). Qualitative and quantitative analysis of phytochemicals of *Taraxacum officinale*. *Wudpecker Journal of Pharmacy and Pharmacology*. 2(1): 001-005.

18. Nguyen, Q. and Eun, J. (2011). *Journal of Medicinal Plants Research*, 5(13): 2798-2811.
19. Oksana, S., Marian, B., Mahendra, R. and Bo, S.H. (2012). Plant phenolic compounds for food, pharmaceutical and cosmetics production. *Journal of Medicinal Plants Research*, 6(13): 2526-2539.
20. Patil, U.H. and Gaikwad, D.K. (2010). Phytochemical evaluation and bactericidal potential of *Terminalia arjuna* stem bark. *Int J Pharm Sci Res.*, 2(3): 614-619.
21. Patino, L.O.J., Prieto, R.J.A. and Cuca, S.L.E. (2012). *Zanthoxylum Genus as Potential Source of Bioactive Compounds, Bioactive Compounds in Phytomedicine*, Prof. Iraj Rasooli (Ed.), ISBN: 978-953-307-805-2, InTech, Available from: <http://www.intechopen.com/books/bioactive-compounds-in-phytomedicine/zanthoxylum-genus-as-potential-source-of-bioactive-compounds>.
22. Pise, N., Jena, K., Maharana, D., Gaikwad, D. and Jagtap, T. (2010). Free radical scavenging potential, reducing power, phenolic and biochemical constituents of *Porphyra* species from India. *J. Algal Biomass Utiln.*, 1(3): 29-42.
23. Prabhash, T., Alagendran, S., ArulBalachandran, D. and Anthonisamy, A. (2014). Assessment of Bioactive compound in *Zanthoxylum rhetsa* (Roxb)-A rare medicinal herb. *Int. J. Cur. Tr. Res.*, 3 (2): 77-83.
24. Prieto, J.A., Patino, O.J., Delgado, W.A., Moreno, J.P. and Cuca, L.E. (2011). Chemical composition, Insecticidal, and Antifungal activities of fruit essential oils of three Colombian *Zanthoxylum* Species. *Chilean Journal of Agricultural Research*, 71(1): 73-82.
25. Qureshi, H., Arshad, M. and Bibi, Y. (2014). Toxicity assessment and phytochemical analysis of *Broussonetia papyrifera* and *Lantana camara*: Two notorious invasive plant species. *J. Bio. & Env. Sci.*, 5(2): 508-517.
26. Rangari, V.D. (2002). *Pharmacognosy and phytochemistry*. Nasik: Carrier Publication, pp: 132.
27. Reddy, L.J. and Jose, B. (2011). Statistical analysis of the antibacterial activity of *Zanthoxylum rhetsa* seed essential oil, *Journal of Chemical and Pharmaceutical Research*, 3(1): 440-444.
28. Sangeetha, C. and Baskar, P. (2015). Allelopathy in weed management-A critical Review. *African Journal of Plant Science*, 10(9):1004-1015.
29. Santhanam, R.K. et al., (2013). Photoprotective properties of *Zanthoxylum rhetsa*: An in vitro analysis. *Journal of Chemical and Pharmaceutical Research*, 5(12): 1512-1520.
30. Shankar, R. and Rawat, M.S. (2013). Conservation and cultivation of threatened and high valued medicinal plants in North East India. *International Journal of Biodiversity and Conservation*, 5(9): 584-591.
31. Singh, H. (2006). Prospects and Challenges for Harnessing Opportunities in Medicinal Plants Sector in India. 2/2 Law, Environment and Development Journal, pp:196, available at <http://www.lead-journal.org/content/06196.pdf>.
32. Singleton, V.L., Orthofer, R. and Lamuela-Raventos, R.M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods Enzymol.*, 299: 152-178.
33. Sofowra, A. (1993). *Medicinal Plants And traditional Medicine in Africa*. Spectrum Books Ltd., Ibadan, Nigeria, pp: 191-289.
34. Trease, G.E. and Evans, W.C. (1989). *Pharmacognosy*, 11<sup>th</sup> Edn. Bailliere Tindall, London, pp: 45-50.
35. Verma, N. and Khosa, R.L. (2012). Hepatoprotective Effect of *Zanthoxylum armatum* DC, *Bioactive Compounds in Phytomedicine*, Prof. Iraj Rasooli (Ed.), ISBN: 978-953-307-805-2, InTech, Available from: <http://www.intechopen.com/books/bioactive-compounds-in-phytomedicine/hepatoprotective-effect-of-zanthoxylum-armatum-dc>.

36. Vidyamadhavi, K., Chandrashekhar, G.J., Manjunath, H.M., Nivya, M.T., Anand, D. and Raju, N.G. (2014). Evaluation of In-Vitro Antioxidant, Anti-inflammatory Properties of Aerial Parts of *Zanthoxylum rhesta*. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 5(5): 997-1003.
37. Wadood, A., Ghufuran, M., Jamal, S.B., Naeem, M., Khan, A., et al. (2013). Phytochemical Analysis of Medicinal Plants Occurring in Local Area of Mardan. *Biochem Anal Biochem.*, 2:144. doi: 10.4172/2161-1009.1000144.
38. World Health Organization, (1999). *Consultation Meeting on Traditional Medicine and Modern Medicine: Harmonizing the Two Approaches*. Geneva, World Health Organization, 1999 (document reference (WP) TM/ICP/TM/001/RB/98–RS/99/GE/32(CHN)).
39. World Health Organization, (2000). *Promoting the Role of Traditional Medicine in Health Systems: a Strategy for the African Region 2001-2010*. Harare, World Health Organization, 2000 (document reference AFR/RC50/Doc.9/R).
40. World Health Organization, (2002). *WHO Traditional Medicine Strategy, 2002-2005*, Geneva, 2002.
41. Yadav, R.N.S. and Agarwala, M. (2011). Phytochemical analysis of some medicinal plants, *Journal of Phytology*, 3(12): 10-14.

